

Vol. 15  
n. 3  
2019

PRS ▶▶▶

# JPD

## JOURNAL of **PLASTIC** *and Pathology* DERMATOLOGY



Official Journal of ISPLAD

Indexed in EMBASE/SCOPUS - ISSN 2035-0686

EDITOR in CHIEF  
Antonino Di Pietro

### Effectiveness of microwaves in the treatment of cellulite: a preliminary study

Antonino Di Pietro, Simona Ferri, Paolo Bonan,  
Alice Verdelli, Silvia Stevan,  
Chiara Tartaglia, Elisabetta Perosino

Periodico trimestrale - Poste Italiane S.p.A. - Spedizione in Abbonamento Postale Aut. MBPVL0-NO/154/A.P./2018 - Stampe Periodiche in Regime Libero - LO/MI

Powered by



# Effectiveness of microwaves in the treatment of cellulite: a preliminary study



ANTONINO DI PIETRO

• Antonino Di Pietro<sup>1</sup> • Simona Ferri<sup>1</sup> • Paolo Bonan<sup>2</sup> • Alice Verdelli<sup>2</sup> • Silvia Stevan<sup>3</sup> • Chiara Tartaglia<sup>4</sup> • Elisabetta Perosino<sup>4</sup>

## SUMMARY

**Background.** Cellulite is a major aesthetic concern affecting approximately 90% of post-puberal women worldwide. Numerous invasive and non-invasive treatments have been proposed, with variable efficacy. The objective of this study was to investigate the effectiveness of a new microwaves platform, ONDA Coolwaves® (DEKA, Florence, Italy), in the treatment of cellulite in the areas of the buttocks and posterior thighs.

**Material and Methods.** Patients with cellulite on the buttocks and posterior thighs were enrolled in two Italian Centers. The treatment protocol consisted of 4 sessions of ONDA Coolwaves® at intervals of 30 days. Clinical evaluations and standardized digital photographs were performed before the first treatment, at 4 and at 8 weeks after the last treatment. To assess the severity of cellulite the cellulite severity scale (CSS) was used.

**Results.** Twenty women were enrolled in this study. At baseline, 95% of the patients showed a moderate-severe CSS. A grade II-III cellulite was found in 85% of the women, while 15% showed a mild cellulite (grade I). Eight weeks after the last treatment, an overall improvement of cellulite grade was found ( $p < 0.05$ ) with 80% of the women showing a mild CSS and 20% of patients showing a moderate CSS index. Totally, 25% of the patients did not have signs of cellulite in the follow-up period, while 75% patients showed a reduction of cellulite to grade I or II.

**Conclusions.** This prospective study confirmed the efficacy of ONDA Coolwaves® (DEKA, Florence, Italy) for the treatment of cellulite on both buttocks and posterior thighs areas, with no detected side effects.

## KEYWORDS

Microwaves, cellulite.

1 Istituto Dermoclinico  
"Vita Cutis", Milan (Italy)

2 Laser Cutaneous Cosmetic  
& Plastic Surgery Unit, Villa  
Donatello Clinic, Florence (Italy)

3 Centro Medico Laser  
Scandicci, Florence (Italy)

4 Studio Medico Perosino  
Associati, Rome (Italy)

## INTRODUCTION

Cellulite is one of the most common lipodystrophy syndromes, which affects 80-90% of post-adolescent women [1]. Firstly described in the early 20<sup>th</sup> century, it has been referred to by several synonyms, including gynoid lipodystrophy, nodular liposclerosis, edematofibrosclerotic panniculopathy, panniculosis, adiposis edematosa, dermopanniculosis deformans, and status protrusus cutis [2-4].

Clinically, cellulite shows off with a surface relief alteration resulting in depression and raised areas leading to an irregular appearance, such as dimpled, orange peel, cottage cheese, or mattress-like appearance of the area of interest, typically located over the thighs and buttocks but also on the legs, arms and abdomen [5]. Changes occurring in the course of cellulite formation regard fibrosis and sclerosis of the tissues, which may result in a wavy course of the skin surface with palpable multiple nodules. Many studies confirmed that cellulite depressions are associated with a significant increase in the presence and thickness of underlying subcutaneous fibrous septa. In cases of significant advancement of changes, nodules and sclerotic lesions are accompanied by soreness, dryness and thinning of the skin covering the changed area [6-9].

Women are most commonly affected by cellulite due to the structure and anatomy of their subcutaneous septa compared to men [10]. Particularly, in women, fibrous septa are usually oriented perpendicularly to the skin surface, creating large, rectangular fat lobules. Fibrosis leads to the retraction

of the septa, pulling the skin down and creating depressions or dimples. The raised areas result from projection of underlying fat lobules into the dermis. In men, the fibrous septa are organized in a crisscross pattern which create smaller subcutaneous fat lobules, reducing the risk of developing cellulite dimples [10].

The etiopathogenesis of cellulite is either multifactorial and unclear, but it is known to incorporate genetic, vascular, hormonal and environmental factors [11]. Most of the current data indicates that either endothelial dysfunction and lymphatic disorders play an important role in the etiology of this phenomenon, causing interstitial matrix alterations and structural changes in subcutaneous adipose tissue [12-15].

Inflammatory factors could play a role in the fibrous septal development too [13]. Significant decreases in the subcutaneous expression of adiponectin, an adipocyte-derived hormone with anti-inflammatory, antifibrotic, and vasodilatory functions, may play a role in cellulite pathogenesis [1,16]. Changes in the subcutaneous tissue could also be worsened by age, due to an increased skin laxity [17]. Moreover, female sex hormones may play a significant role in the development of lipodystrophy. Particularly, the exacerbation and progression of cellulite seems to correlate with pregnancy-related hyperestrogenism, the use of birth control pills or hormone replacement therapy in postmenopausal women [11]. Alcohol consumption stimulates lipogenesis and causes body dehydration, resulting in the excessive and improper storage of fat [18]. Smoking significantly

contributes to a greater amount of free radicals production in the body, and to contract the small blood vessels, responsible for local micro-circulatory impairments [19]. Finally, lifestyle factors such as high carbohydrate diets, and a sedentary life may lead to an increase in total body fat content, which enhances the appearance of cellulite [1].

To assess the severity of cellulite, Nürnberger and Müller [4] first proposed a clinical grading scale for cellulite, including three grades:

- 1. Grade I= the skin of the affected area is smooth while the subject is standing or lying, but the alterations to the skin surface can be seen by pinching the skin or with muscle contraction (visible changes with skin clamping or muscle contraction);
- 2. Grade II= the orange peel aspect of the skin or mattress appearance is evident when standing, without the use of any manipulation of the skin pinching or muscle contraction (visible without manipulation);
- 3. Grade III= the alterations described in grade or stage II, are present together with raised areas and nodules (visible changes associated with nodules).

Recently, Hexsel et al. [20] validated a standardized and objective method of grading cellulite, the *cellulite severity scale* (CSS). It is based on 5 key morphologic aspects of cellulite, including the 1) number of depressions; 2) depth of depressions; 3) clinical morphology; 4) extent of skin laxity, flaccidity, or sagging; and 5) Nürnberger-Müller classification grade. Each variable is graded from 0 to 3, leading to overall grades of mild (1-5), moderate (6-10), and severe (11-15).

During the years, many treatments have been proposed to improve cellulite, including topical therapy, injectables (eg, chemical septolysis with collagenase), lymphatic or vacuum-assisted massage, acoustic wave therapy, light therapy, external non-invasive lasers, and radiofrequency (RF) devices [21,22]. Unfortunately, unpredictable efficacy and the potential for only short-term improvement despite numerous treatment sessions limit their popularity.

Herein, we describe a non-invasive treatment for women with cellulite, evaluating the safety and effectiveness of a new technique that utilizes microwaves for the reduction of the visible appearance of cellulite, the ONDA Coolwaves® system (DEKA, Florence, Italy).

## Materials and methods

This prospective study was performed in two Italian Centers (Istituto Dermoclinico Vita Cutis e Studio Medico Perosino Associati) from September 2018 to June 2019.

Female patients with gluteal and posterior thighs cellulite, aged > 18 years with a body mass index (BMI) <30 were enrolled.

### Exclusion Criteria:

- Volunteers undergoing other cosmetic treatment in the last 6 months prior to the study;
- Pregnant or lactating patients;
- Patients having both present or past history of cardiovascular diseases, cancers, metabolic disorders, immunosuppression.

After being informed about the aim of the study, patients gave their

written informed consent. Moreover, all the patients gave consent to the use of their photographs for scientific purposes. All the enrolled volunteers were properly informed about the treatment steps and procedures to be performed. The participants were not submitted to any food restriction and were asked to maintain their usual daily activities.

The treatment protocol consisted of 4 sessions of ONDA Coolwaves® (DEKA, Florence, Italy) spaced 30 days apart each other. The ONDA Coolwaves® is a microwaves platform utilizing two handpieces which allow a selective concentration of the specific microwave dielectric heating in both deep and superficial subcutaneous tissues. An effective contact cooling system is implemented to avoid rebounding thermal damage to the superficial layers during the microwaves transmission.

Clinical evaluations were performed before the first treatment (at baseline), at 4 and at 8 weeks after the last treatment. Cellulite grade was determined by clinical inspection of the patient's skin. To assess cellulitis severity, the CSS was used. The treated areas were inspected and constantly monitored during all the sessions. Standardized digital photographs with a digital camera (Reflex Nikon D800) were also performed at baseline, at 4 and at 8 weeks after the last treatment.

Patients comfort and satisfaction were evaluated using a five-point Likert scale questionnaire (0 = worse; 1 = little satisfaction or not satisfied; 2 = fairly satisfied; 3 = satisfied; and 4 = very satisfied).

All the clinical and anthropometric data, including age and BMI

of the patients were recorded and put into a database. The Wilcoxon signed rank test was used for statistical analysis, with an alpha level of 5% ( $p < 0.05$ ) being considered significant. Data were represented as means  $\pm$  standard deviation (SD).

## Results


### → Demographic data

A total of 20 women took part in the study and completed all the study visits. Mean age ( $\pm$  SD) at baseline was  $37.0 \pm 12.1$  years (range, 19–56 years) and mean was BMI  $21.7 \pm 2.2$  kg/m<sup>2</sup> (range, 19–28 kg/m<sup>2</sup>).


### → Cellulite Severity Scale

At baseline, the mean CSS ( $\pm$  SD) was  $9.2 \pm 2.3$ . 75% out of the 20 patients (n=15) showed a moderate CSS, followed by 20% of women with a severe CSS (n=4) and only 5% (n=1) patients with a mild CSS (Figure 1). According to Nürnberg and Müller classification, 65% (n=13) of the women rated as grade II and 20% (n=4) as grade III cellulite; 15% of the women (n=3) showed a grade I cellulite (Figure 2). Moderate-severe skin depressions as well as medium-severe depth depressions were present in 75% (n=15) out of the 20 patients. Concerning skin laxity, 65% (n=13) of the women showed slight laxity, while 35% (n=5) presented a moderate form.

At follow-up, 8 weeks after the last treatment, all the parameters analyzed according to CSS, were improved ( $p < 0.05$ ). Mean CSS was  $4.2 \pm 1.9$ . Totally, 80% (n=16) of the women showed a mild CSS while 20% (n=4) of the patients presented a mo-



**Figure 1.** Cellulite severity scale (CSS) at baseline and 8 weeks after the last session of ONDA Coolwaves® (DEKA, Florence, Italy).



**Figure 2.** Nürnberger-Müller classification (grade 0-3) at baseline and 8 weeks after the last session of ONDA Coolwaves® (DEKA, Florence, Italy).

derate CSS index (**Figure 1**). None of the women showed severe CSS. An overall improvement of cellulite grade, according to Nürnberger and Müller classification, was shown ( $p < 0.05$ ) with 25% ( $n=5$ ) of the patients with no signs of cellulite, while 60% ( $n=12$ ) out of the 20 patients showed a grade I cellulitis and 15% ( $n=3$ ) a grade II cellulite (**Figure 2**).

Only 10% of the patients ( $n=2$ ) showed moderate depressions, while 65% ( $n=13$ ) and 15% ( $n=5$ ) patients showed mild or did not show depressions, respectively. Moreover, 20% ( $n=4$ ) out of 20 patients presented severe depth depressions, while 65% women ( $n=13$ ) showed medium depth depressions. Skin laxity was absent in 65% ( $n=13$ ) of patients and mild ( $n=7$ ) in 35% of women.

Among volunteers, no significant statistical BMI variation was found.

#### → Side effects

None of the volunteers reported either pain or other side effects such as

post-operative ecchymosis, oedema, blistering, hardness, neurovascular and local tissue injury (e.g. temporary loss of cutaneous sensory), swelling or nodules during either the treatment nor in the follow-up period.

#### → Patient Satisfaction Index

Seventeen out of the twenty patients reported a significant improvement of cellulite.

On a scale of 0-4, the patients agreed (3.8) that they were satisfied with the results of the treatment. Three patients reported a mild satisfaction index because cellulite was improved but not completely disappeared. Patients also found an improvement in skin laxity.

## Discussion

Cellulite is a condition that affects psychological aspect in majority of women. Many therapeutic options have been

**Figure 3.**  
Abdomen cellulite of a 42 female patient before (a) and after (b) 1 month of treatment with ONDA Coolwaves® (clinical photography).



**Figure 4.**  
Buttocks and thighs cellulite of a 38 female patient before (a) and after (b) 4 months of treatments with ONDA Coolwaves® (clinical photography).



developed to overcome this condition, including topical treatments, non-invasive energy-based devices, and minimally invasive interventions [1,5,12,13,22,23,24].

RF techniques are one of the most commonly treatment for cellulite. Many companies propose RF systems (100 kHz–40 MHz) as valid and reliable alternatives to other technologies aimed at reaching the same clinical results [25]. RF is an electromagnetic energy that works by heating the tissues. When applied to tissues, RF generates oscillating magnetic fields that move electrically charged particles, producing heat in the tissues, and the amount of heat produced depends on the resistance (bio-impedance) of the target tissue [26–28]. The heat that is delivered to the subcutaneous layer is presumably absorbed by adipocytes till, to supposedly induce the breaking down of fat cells through the membrane's lysis [29,30]. Subsequently, a wound-healing process such as collagen neo-synthesis would therefore improve various tissue characteristics [30,31]. By elevating the tissue temperature at the target area, some collagen denaturation, remodelling and new collagen generation are stimulated. According to the literature, RF also favours a decrease in thickness and fat accumulation in adipocytes, consequently reducing venous and lymphatic fluid retention caused by hypodermal tissue compressing vessels and nerve endings [32].

RF generates heat in different tissues by transforming energy through three basic mechanisms from electromagnetic field. These mechanisms include (i) orientation of electric dipoles that already exists in the atoms and molecules in the tissue; (ii) polarization of atoms and mole-

cules to produce dipole moments; and (iii) displacement of conduction electrons and ions in the tissue [33].

RF devices deliver thermal energy via electrode. Electro-thermal energy production with this modality could be monopolar (single electrode with return electrode), unipolar (single electrode with no return), bipolar (2 electrodes), or multipolar (3 or more electrodes) [24,34]. Recently, contactless RF devices have also been developed. RF has been used alone or in combination with other treatments, such as infrared light and mechanical manipulation [24,37,38].

Current studies focusing on RF in the treatment of cellulite were reported in **Table 1** [27–29, 39–54]. Among the RF devices, combination therapy is the most investigated technique, with ten published scientific papers. Six randomized controlled trials (RCTs) were also available.

In a recent review by Vale et al., [26] authors reported all the current literature on RF in the treatment of cellulite. Generally, authors showed the efficacy of high-energy RF and RF combination therapy for treating cellulite; however, authors underlined that the methodological quality of these studies was low and that statistical analyses was not performed in all the studies. Moreover, not all the studies used a standardized method to objectively evaluate the results of the treatment, thus making difficult to confirm their results.

Despite multiple RF devices that attempt to treat cellulite, no procedures have been proven to be successful in the long term [32]. As the demand for non-invasive and long-lasting treatments to reduce cellulite has been growing up, new medical devices have been developed accordingly. Recently, a new microwave techno-

logy has been setup [33]. Microwaves are part of the RF spectrum; they are electromagnetic waves with a frequency range between 1 and 300 GHz [55]. In a previous study by Bonan et al., [56] ONDA Coolwaves<sup>®</sup>, (DEKA, Florence, Italy) a new microwave device, has been successfully used for the treatment of localized subcutaneous adiposities. The ONDA system handpieces generate waves at 2.45 GHz which interact with biological molecules and generates localized, controlled heat absorbed by selected biological targets, such as fat, through a biophysical process called “dielectric heating”. Authors hypothesized that microwaves could act directly on the adipose tissue, sparing the dermo-epidermal layers, promoting adipocyte cells heating, and, consequently, leading to macrophage-mediated adipocytes lysis. Interestingly, authors found an improvement in the patients' skin tone and texture. Authors hypothesize that this improvement could be the result of the action of microwaves on collagen septa. Controlled hyperthermia could cause solubilisation of the deeper collagen fibers and activate fibroblasts, remodelling collagen fibers. On these basis, we aimed at evaluating if microwaves could also improve cellulite. The objective of our study was to verify the effectiveness and safety of ONDA Coolwaves<sup>®</sup> in the treatment of cellulite and its long-term efficacy.

To assess the severity of cellulite in a standardized and objective way, the CSS was used. We found a significant improvement of cellulite with a significant reduction of number and depth of depressions in all the patients. The mean total CSS score improved from moderate-to-severe at baseline to mild 8 weeks after the protocol treatment with ONDA Coolwaves<sup>®</sup>.

No side effects were reported, neither after the single sessions of treatment, nor in the follow-up period, making the treatment safe and well tolerated by all the patients. Our results also showed a long-term efficacy of the treatment, since benefits were kept up to 8 weeks after the last session of treatment (totally 6 months). Interestingly, during the treatment period, there were no significant changes in the BMI, confirming previous data [56].

One of the limits of our study could be the small number of enrolled patients. Therefore, further studies with a larger number of patients will be needed to more fully characterize the clinical potential of microwaves and their mechanism of action.

## Conclusions

This prospective study confirmed that ONDA Coolwaves® (DEKA, Florence, Italy) treatment is a safe and effective treatment for cellulite of the buttocks and posterior thighs. Moreover, the evaluation of parameters through the CSS before and 8 weeks after the last treatment, proved a long-lasting effect of ONDA Coolwaves®.

*Conflict of interest: none*

*Financial disclosure: none*

**Table 1.**  
Overview of studies  
evaluating the efficacy of  
radiofrequency in cellulite.

| References            | Year | Design | Therapy       | N° patients | Statistical analysis |
|-----------------------|------|--------|---------------|-------------|----------------------|
| Sadick NS             | 2004 | OS     | Unipolar RF   | 35          | no                   |
| Alster TS             | 2005 | OS     | Combination   | 20          | no                   |
| Emilia del Pino M     | 2006 | OS     | Combination   | 26          | no                   |
| Kulick M              | 2006 | OS     | Combination   | 16          | no                   |
| Wanitphakdeedecha R   | 2006 | OS     | Combination   | 12          | no                   |
| Nootheti PK           | 2006 | RCT    | Combination   | 20          | yes                  |
| Sadick N              | 2007 | OS     | Combination   | 16          | no                   |
| Alexiades-Armenakas M | 2008 | RCT    | Unipolar RF   | 10          | yes                  |
| Goldberg DJ           | 2008 | OS     | Unipolar RF   | 30          | no                   |
| Romero C              | 2008 | RCT    | Combination   | 10          | yes                  |
| Van der Lugt C        | 2009 | OS     | Bipolar RF    | 50          | no                   |
| Manuskiatti W         | 2009 | OS     | Low-level RF  | 37          | yes                  |
| Kaplan H              | 2009 | OS     | Low-level RF  | 12          | no                   |
| Boisnic S             | 2010 | RCT    | Low-level RF  | 24          | yes                  |
| Levenberg A           | 2010 | OS     | Low-level RF  | 37          | yes                  |
| Hexsel DM             | 2011 | OS     | Combination   | 11          | yes                  |
| Mlosek RK             | 2012 | RCT    | Low-level RF  | 45          | yes                  |
| McDaniel D            | 2015 | OS     | High-level RF | 30          | yes                  |
| Hayre N               | 2016 | RCT    | High-level RF | 36          | yes                  |
| Suh DH                | 2017 | OS     | High-level RF | 12          | no                   |
| Wanitphakdeedecha R   | 2017 | OS     | Combination   | 25          | yes                  |
| Firtz K               | 2018 | OS     | Combination   | 30          | yes                  |

OS= observational study, RCT= randomized controlled study, RF= radiofrequency.



## REFERENCES

1. Friedmann DP, Vick GL, Mishra V. Cellulite: a review with a focus on subcision. *Clin Cosmet Investig Dermatol*. 2017 Jan 7;10:17-23.
2. Rossi AB, Vergnanini AL. Cellulite: a review. *J Eur Acad Dermatol Venereol*. 2000;14: 251–262.
3. Avram MM. Cellulite: a review of its physiology and treatment. *J Cosmet Laser Ther*. 2004; 6:181–185.
4. Nürnberg F, Müller G. So-called cellulite: an invented disease. *J Dermatol Surg Oncol*. 1978; 4:221–229.
5. Rossi AM, Katz BE. A modern approach to the treatment of cellulite. *Dermatol Clin*. 2014;32:51–59.
6. Mirrashed F, Sharp JC, Krause V, Morgan J, et al. Pilot study of dermal and subcutaneous fat structures by MRI in individuals who differ gender, BMI, and cellulite grading. *Ski Res Technol*. 2004;10: 161–168.
7. Meyer PF, Martins NM, Martins FM, Monteiro RA, de Mendonça KMPP. Effects of lymphatic drainage on cellulitis assessed by magnetic resonance. *Brazilian Arch Biol Technol*. 2008;51(Special issue):2 21–224.
8. Hexsel DM, Abreu M, Rodrigues TC, et al. Side-by-side comparison of areas with and without cellulite depressions using magnetic resonance imaging. *Dermatol Surg*. 2009;35: 1471–1477.
9. Omi T, Sato S, Kawana S. Ultrastructural assessment of cellulite morphology: clues to a therapeutic strategy? *Laser Ther*. 2013;22: 131–136.
10. Querleux B, Cornillon C, Jolivet O, et al. Anatomy and physiology of subcutaneous adipose tissue by *in vivo* magnetic resonance imaging and spectroscopy: relationships with sex and presence of cellulite. *Skin Res Technol*. 2002;8: 118–124.
11. Tokarska K, Tokarski S, Wo niacka A, et al. Cellulite: a cosmetic or systemic issue? Contemporary views on the etiopathogenesis of cellulite. *Postepy Dermatol Alergol*. 2018;35: 442–446.
12. Mazioti M. The potential role of endocrine disrupting chemicals in cellulite. *Med Hypotheses*. 2018;116:132–135.
13. Hogan S, Velez MW, Kaminer MS. Updates on the understanding and treatment of cellulite. *Semin Cutan Med Surg*. 2018;37: 242–246.
14. Callaghan DJ Rd, Robinson DM, Kaminer MS. Cellulite: a review of pathogenesis-directed therapy. *Semin Cutan Med Surg*. 2017;36 :179–184.
15. Curri SB. Cellulite and fatty tissue microcirculation. *Cosmet Toilet*. 1993;108:51–58.
16. Emanuele E, Minoretta P, Altabas K, et al. Adiponectin expression in subcutaneous adipose tissue is reduced in women with cellulite. *Int J Dermatol*. 2011;50: 412–416.
17. Casabona G, Pereira G. Microfocused Ultrasound with Visualization and Calcium Hydroxylapatite for Improving Skin Laxity and Cellulite Appearance. *Plast Reconstr Surg Glob Open*. 2017;5:e1388.
18. Steiner JL, Lang CH. Alcohol, Adipose Tissue and Lipid Dysregulation. *Biomolecules*. 2017;7: 16.
19. Stavroulaki A, Pramantiotis G. Cellulite, smoking and angiotensin converting enzyme (ACE) gene insertion/deletion polymorphism. *J Eur Acad Dermatol Venereol*. 2011;25: 1116–1117.
20. Hexsel DM, Dal’Forno T, Hexsel CL. A validated photonumeric cellulite severity scale. *J Eur Acad Dermatol Venereol*. 2009;23: 523–8.
21. Luebberding S, Krueger N, Sadick NS. Cellulite: an evidence-based review. *Am J Clin Dermatol*. 2015;16:243–256.
22. Sadick N. Treatment for cellulite. *Int J Womens Dermatol*. 2019; 5: 68–72.
23. Zerini I, Sisti A, Cuomo R, et al. Cellulite treatment: a comprehensive literature review. *J Cosmet Dermatol*. 2015;14:224–40.
24. Friedmann DP, Vick GL, Mishra V. Cellulite: a review with a focus on subcision. *Clin Cosmet Investig Dermatol*. 2017;10:17–23.
25. Belenky I, Margulis A, Elman M, Bar-Yosef U, Paun SD. Exploring Channeling Optimized Radiofrequency Energy: a Review of Radiofrequency History and Applications in Esthetic Fields. *Adv Ther*. 2012; 29:249–266.
26. Vale AL, Pereira AS, Morais A, Noites A, Mendonça AC, Martins Pinto J, Vilarinho R, Carvalho P. Effects of radiofrequency on adipose tissue: A systematic review with meta-analysis. *J Cosmet Dermatol*. 2018;17:703–711.
27. Alexiades-Armenakas M, Dover JS, Arndt KA. Unipolar radiofrequency treatment to improve the appearance of cellulite. *J Cosmet Laser Ther*. 2008;10:148–153.
28. Kaplan H, Gat A. Clinical and histopathological results following TriPollar radiofrequency skin treatments. *J Cosmet Laser Ther*. 2009; 11:78–84.
29. Van der Lugt C, Romero C, Ancona D, et al. A multicenter study of cellulite treatment with a variable emission radio frequency system. *Dermatol Ther*. 2009; 22:74–84.
30. Bojsnic S, Branchet M-C, Birnstiel O, et al. Clinical and histopathological study of the TriPollar home-use device for body treatments. *Eur J Dermatol*. 2010;20:367–72.

31. Fisher GH, Jacobson LG, Bernstein LJ, et al. Nonablative radiofrequency treatment of facial laxity. *Dermatol Surg.* 2005;31:1237–41.
32. Da Silva RMV, Arend Barichello P, Lima Medeiros M. Effect of Capacitive Radiofrequency on the Fibrosis of Patients with Cellulite. *Dermatol Res Pract* 2013; 2013:715829.
33. Alizadeh Z, Halabchi F, Mazaheri R, et al. Review of the Mechanisms and Effects of Noninvasive Body Contouring Devices on Cellulite and Subcutaneous Fat. *Int J Endocrinol Metab.* 2016;14:e36727.
34. Peterson JD, Goldman MP. Laser, light, and energy devices for cellulite and lipodystrophy. *Clin Plast Surg.* 2011;38:463–474.
35. McDaniel D, Samkova P. Evaluation of the safety and efficacy of a non-contact radiofrequency device for the improvement in contour and circumferential reduction of the inner and outer thigh. *J Drugs Dermatol.* 2015;14:1422-1424.
36. Hayre N, Palm M, Jenkin P. A clinical evaluation of a next generation, non-invasive, selective radiofrequency, hands-free, body-shaping device. *J Drugs Dermatol.* 2016;15:1557-1561.
37. Krueger N, Sadick NS. New-generation radiofrequency technology. *Cutis.* 2013;91:39-46.
38. Foster, K. R., Ziskin, M. C., & Balzano, Q. Thermal response of human skin to microwave energy: A critical review. *Health Physics,* 206:111; 528–541.
39. Sadick NS, Mulholland RS. A prospective clinical study to evaluate the efficacy and safety of cellulite treatment using the combination of optical and RF energies for subcutaneous tissue heating. *J Cosmet Laser Ther.* 2004;6:187–90.
40. Alster TS, Tanzi EL. Cellulite treatment using a novel combination radiofrequency, infrared light, and mechanical tissue manipulation device. *J Cosmet Laser Ther.* 2005;7:81–5.
41. Emilia del Pino M, RosadoRH, AzuelaA, et al. Effect of controlled volumetric tissue heating with radiofrequency on cellulite and the subcutaneous tissue of the buttocks and thighs. *J Drugs Dermatol.* 2006;5:714-722.
42. Kulick M. Evaluation of the combination of radio frequency, infrared energy and mechanical rollers with suction to improve skin surface irregularities (cellulite) in a limited treatment area. *J Cosmet Laser Ther.* 2006;8:185–90.
43. Nootheti PK, Magpantay A, Yosowitz G, et al. A single center, randomized, comparative, prospective clinical study to determine the efficacy of the VelaSmooth system versus the Triactive system for the treatment of cellulite. *Lasers Surg Med.* 2006;38:908–12.
44. Wanitphakdeedecha R, Manuskiatti W. Treatment of cellulite with a bipolar radiofrequency, infrared heat, and pulsatile suction device: a pilot study. *J Cosmet Dermatol.* 2006;5:284–8.
45. Sadick N, Magro C. A study evaluating the safety and efficacy of the VelaSmooth system in the treatment of cellulite. *J Cosmet Laser Ther.* 2007;9:15–20.
46. Goldberg DJ, Fazeli A, Berlin AL. Clinical, laboratory, and MRI analysis of cellulite treatment with a unipolar radiofrequency device. *Dermatol Surg.* 2008;34:204–9.
47. Romero C, Caballero N, Herrero M, et al. Effects of cellulite treatment with RF, IR light, mechanical massage and suction treating one buttock with the contralateral as a control. *J Cosmet Laser Ther.* 2008;10:193–201.
48. Manuskiatti W, Wachirakaphan C, Lektrakul N, et al. Circumference reduction and cellulite treatment with a TriPollar radiofrequency device: a pilot study. *J Eur Acad Dermatol Venereol.* 2009;23:820–7.
49. Levenberg A. Clinical experience with a TriPollar radiofrequency system for facial and body aesthetic treatments. *Eur J Dermatol.* 2010;20:615-619
50. Hexsel DM, Siega C, Schilling-Souza J, Porto MD, Rodrigues TC. A bipolar radiofrequency, infrared, vacuum and mechanical massage device for treatment of cellulite: a pilot study. *J Cosmet Laser Ther.* 2011;13:297–302.
51. Mlosek RK, Wozniak W, Malinowska S, Lewandowski M, Nowicka A. The effectiveness of anticellulite treatment using tripolar radiofrequency monitored by classic and high-frequency ultrasound. *J Eur Acad Dermatol Venereol.* 2012;26:696-703.
52. Suh DH, Kim CM, Lee SJ, Kim H, Yeom SK, Ryu HJ. Safety and efficacy of a non-contact radiofrequency device for body contouring in Asians. *J Cosmet Laser Ther.* 2017;19:89-92.
53. Wanitphakdeedecha R, Sathaworawong A, Manuskiatti W, et al. Efficacy of multipolar radiofrequency with pulsed magnetic field therapy for the treatment of abdominal cellulite. *J Cosmet Laser Ther.* 2017;19:205-209.
54. Fritz K, Salavastru C, Gyurova M. Reduction of abdominal skin laxity in women postvaginal delivery using the synergistic emission of radiofrequency and targeted pressure energies. *J Cosmet Dermatol.* 2018;17:766-769.
55. Beasley KL, Weiss RA. Radiofrequency in cosmetic dermatology. *Dermatol Clin.* 2014;32:79-90.
56. Bonan P, Marini L, Lotti T. Microwaves in body sculpting: A prospective study. *Dermatol Ther.* 2019;32:e12782.